

Dependence on Ionic Strength of Formation Constants, Protonation, and Complexation of Nitrilotriacetic Acid with Tungsten(VI) in Sodium Perchlorate Aqueous Solution

Kavosh Majlesi,^{*†} Karim Zare,^{†‡} and Faranak Teimouri[†]

Chemistry Department, Islamic Azad University, Science & Research Campus, Tehran, Hesarak, Iran, and Chemistry Department, Shahid Beheshti University, Tehran, Evin, Iran

By use of potentiometric and spectrophotometric techniques, the complexation of tungsten(VI) with nitrilotriacetic acid (NTA) has been carried out in aqueous solution for pH = 7.5 at 25 °C and different ionic strengths ranging from (0.1 to 1.0) mol dm⁻³ (NaClO₄). The composition of the complex was determined by the continuous variations method. It was shown that tungsten(VI) forms a mononuclear complex with NTA of the type (WO₃L³⁻) at pH = 7.5. The dependence of the protonation of the aforementioned ligand and the stability constant of the complex on ionic strength is described by a Debye–Huckel-type equation. Finally the results have been compared with the previously reported data.

Introduction

Desulfovibrio gigas formate dehydrogenase is the first representative of a tungsten-containing enzyme from a mesophile that has been structurally characterized.¹ It is a heterodimer of 110 and 24 kDa subunits. The existence of a universal pterin dithiolene cofactor ligand for the molybdenum and tungsten oxotransferases supports a biological significance of the fundamental chemistry of mono- and bis(dithiolene) complexes of these elements.² Tungsten(VI) also forms complexes with tryptophan³ and porphyrin.⁴ By pursuing our systematic study on complexation of molybdenum(VI) with nitrilotriacetic acid (NTA), we thought it interesting to investigate the protonation and complexation of NTA with tungsten(VI) at different ionic strengths of sodium perchlorate.

Considerable studies have been performed on the ionic strength dependence of the formation constants of some weak complexes of alkali and alkaline earth metals with some organic and inorganic ligands,^{5–13} and little work has been carried for more stable complexes of α -amino acids with transition-metal ions,^{14–17} but according to literature, no work has been reported on the ionic strength dependence of formation constants, protonation, and complexation of tungsten(VI) with NTA.

Marcu¹⁸ studied the radiochromatographic and electrochromatographic of sodium tungstate solutions under the action of NTA. Studies on the structural and bonding characteristics of various Mo(VI)–aminopolycarboxylic acid complexes has led to the evaluation of stability constants from proton nuclear magnetic resonance (NMR) data.^{19,20} On the basis of these studies, Kula²¹ determined the stability constants of W(VI)–NTA complexes by potentiometric techniques. Collin²² reported the stability constant of this system at 25 °C and at an ionic strength of 0.5 mol dm⁻³ NaClO₄. Zare²³ studied this system and evaluated its stability constant at 25 °C and at an ionic strength of 3

mol dm⁻³ NaClO₄ using the potentiometric technique. Anderge²⁴ has made a critical survey of stability constants of NTA complexes.

In the present work, we have studied the complexes of W(VI) with NTA in an ionic strength range of (0.1 to 1.0) mol dm⁻³ sodium perchlorate at 25 °C. A simple Debye–Huckel-type equation was established for the dependence of formation constants on ionic strength. This equation makes it possible to estimate a stability constant at a fixed ionic strength when its value is known at another ionic media in the range (0.1 < I < 1.0) mol dm⁻³.

Experimental Section

Reagents. Sodium perchlorate, perchloric acid, sodium hydroxide, sodium tungstate, and NTA were obtained from E. Merck as analytical reagent grade materials and were used without further purification. Dilute perchloric acid solution was standardized against KHCO₃. In all experiments, double-distilled water with specific conductance equal to (1.3 ± 0.1) $\mu\Omega^{-1}$ cm⁻¹ have been used.

Measurements

A Horiba pH meter, D-14, was used for pH measurements. The pH meter has a sensitivity of 0.01 units. The hydrogen ion concentration was measured with a Horiba combination electrode, model S8720. A 0.01 mol dm⁻³ perchloric acid solution containing 0.09 mol dm⁻³ sodium perchlorate (for adjusting the ionic strength to 0.1 mol dm⁻³) was employed as a standard solution of hydrogen ion concentration. The same procedure was performed for the other ionic strengths.²³ The calibration has been done for the whole pH (pH = -log[H⁺]) range used. Spectrophotometric measurements were performed on a UV–vis Shimadzu 2101 spectrophotometer with an Acer Mate 486 SX/250 computer using thermostated, matched 10-mm quartz cells.

For each experiment two solutions of W(VI) + NTA were prepared with the same concentration, but the ionic strength of the first was maintained with sodium perchlorate and that of the second with sodium hydroxide or

* Author to whom correspondence should be addressed. E-mail: kavoshmajlesi@hotmail.com.

† Islamic Azad University.

‡ Shahid Beheshti University.




Table 1. Potentiometric Data for the Titration of NTA with Sodium Hydroxide at $I = 0.1 \text{ mol dm}^{-3}$ of NaClO_4 and 25°C

V/mL	pH	V/mL	pH	V/mL	pH	V/mL	pH
0.00	2.49	1.64	3.69	2.00	8.86	3.66	11.04
0.01	2.50	1.67	3.79	2.02	8.93	3.72	11.11
0.02	2.50	1.70	3.93	2.04	8.98	3.78	11.17
0.04	2.51	1.72	4.06	2.06	9.04	3.85	11.23
0.06	2.51	1.74	4.20	2.08	9.09	3.95	11.32
0.09	2.52	1.75	4.32	2.10	9.13	4.05	11.39
0.12	2.53	1.76	4.41	2.12	9.18	4.15	11.45
0.16	2.54	1.77	4.51	2.15	9.25	4.25	11.49
0.20	2.55	1.78	4.68	2.18	9.32	4.40	11.57
0.30	2.58	1.79	5.03	2.21	9.37	4.60	11.65
0.40	2.62	1.80	5.44	2.25	9.45	4.80	11.71
0.45	2.64	1.81	5.78	2.29	9.52	5.10	11.79
0.50	2.65	1.82	6.36	2.33	9.60	5.40	11.85
0.60	2.70	1.83	6.84	2.35	9.63	5.80	11.93
0.70	2.74	1.84	7.20	2.40	9.72	6.20	11.99
0.80	2.79	1.85	7.66	2.45	9.81	6.70	12.04
0.90	2.84	1.86	7.81	2.50	9.89	7.20	12.09
1.00	2.90	1.87	7.96	3.10	10.00	7.70	12.14
1.10	2.96	1.88	8.10	3.16	10.10	8.20	12.18
1.20	3.04	1.89	8.23	3.22	10.22	9.00	12.24
1.30	3.13	1.90	8.34	3.28	10.32	10.00	12.31
1.35	3.18	1.91	8.37	3.34	10.44	11.00	12.35
1.40	3.24	1.93	8.52	3.40	10.55	12.00	12.44
1.45	3.30	1.95	8.65	3.46	10.66	14.00	12.50
1.50	3.38	1.96	8.70	3.52	10.77	16.00	12.55
1.55	3.47	1.97	8.74	3.58	10.87		
1.60	3.57	1.98	8.80	3.64	10.96		

perchloric acid. The pH of the first solution was adjusted with the second one. The second solution consists of metal + ligand + NaOH for increasing pH, while for decreasing pH, the second one consists of metal + ligand + HClO_4 . The absorbance of the first solution was measured after adjusting the pH. In all cases, the procedure was repeated at least three times and the resulting average values and corresponding standard deviations are shown in the text and Tables.

Results and Discussion


The dissociation equilibria of NTA have been studied in different kinds of background electrolytes but there are no reports about the ionic strength dependence of the dissociation constants of NTA.²⁵ The following equilibria were studied



Where L^{3-} represents the fully dissociated NTA anion. Three titrations have been done for each ionic strength, and 106 points have been used for calculations at each ionic strength. The dissociation constants K_1 , K_2 , and K_3 have been determined using potentiometric techniques and

Table 2. Dissociation Constants K_1 , K_2 , and K_3 of NTA at Different Ionic Strengths, I , of NaClO_4

$I/\text{mol dm}^{-3}$	$\log K_1$	$\log K_2$	$\log K_3$	experimental conditions	ref
0.1	1.98 ± 0.02	2.92 ± 0.05	10.00 ± 0.02		this work
0.3	1.83 ± 0.04	2.84 ± 0.03	9.85 ± 0.01		this work
0.5	1.76 ± 0.01	2.79 ± 0.02	9.70 ± 0.03		this work
0.7	1.63 ± 0.02	2.61 ± 0.01	9.55 ± 0.02		this work
1.0	1.55 ± 0.02	2.59 ± 0.04	9.25 ± 0.02		this work
	2.05 ± 0.05	2.63 ± 0.02	9.81 ± 0.10	$I = 0.15 \text{ M}, T = 25^\circ\text{C}$	21
	1.65	2.94	9.17 ± 0.04	$I = 3 \text{ M NaClO}_4, T = 25^\circ\text{C}$	23
			10.33	$T = 20^\circ\text{C}$	29

**Figure 1.** Continuous-variation plots of the absorbances of WO_3L^{3-} , abs, vs the mole fraction of W(VI) , X_w , at 25°C , an ionic strength of 0.1 mol dm^{-3} NaClO_4 , and different wavelengths: A, 260 nm; B, 265 nm; and C, 270 nm.

calculated using Microsoft Excel 2000 program.^{25,26} Potentiometric data are listed in Table 1. Dissociation constants are given in Table 2 together with the values reported in the literature, which are in good agreement with those reported before.

Complexation of Tungsten(VI) with NTA

By use of the continuous variations method, the absorbances of solutions of W(VI) and NTA of total concentration $0.006 \text{ mol dm}^{-3}$ in the UV range (260–270 nm) at a constant pH of 7.5 were determined. When solutions of tungstates are made weakly acidic, polymeric anions are formed, but from more strongly acidic solutions, substances often called tungstic acid are obtained.²⁷ The behavior of the tungstate systems is similar to that of the molybdate systems. Again the degree of aggregation in solution increases as the pH is lowered, and numerous tungstates have been crystallized from the solutions at different pHs.²⁷ Tungsten(VI) and the complex both have absorbances at the specified wavelengths, so the absorbance of the tungsten should be eliminated from the total absorbance (A_{obs}). The observed absorbances were corrected from eq 4 and are summarized in Table 3 and plotted in Figure 1

$$A_c = A_{\text{obs}} - \epsilon_0[\text{W}] \quad (4)$$

A_c , A_{obs} , and ϵ_0 are the absorbance of the complex, the observed absorbance, and the molar absorptivity of W , respectively. ϵ_0 values were calculated at the mole fraction of W equal to 1 and are shown in Table 4. In Figure 1, a maximum at a mole fraction of W equal to 0.5 was obtained, indicating a 1:1 complex. The molar absorptivity of the complex, ϵ_1 , was calculated from the linear part of the aforementioned plot at low mole fraction of W , where essentially all the metal ions were in the form of a complex, and is listed in Table 4. At the maximum point of the plot,

Table 3. Continuous-Variation Data for the W(VI) + NTA System at pH 7.5, an Ionic Strength of 0.1 M NaClO₄, and Different Wavelengths

mole fraction of W(VI)	A^a		
	260 nm	265 nm	270 nm
0.00	0.000	0.000	0.000
0.10	0.120	0.112	0.100
0.20	0.194	0.200	0.198
0.30	0.200	0.230	0.240
0.40	0.213	0.254	0.255
0.50	0.307	0.356	0.356
0.60	0.243	0.292	0.292
0.80	0.143	0.212	0.182
1.00	0.000	0.000	0.000

^a The corrected absorbance of $[W(VI)] + [NTA] = 0.006 \text{ mol dm}^{-3}$.


Table 4. Molar Absorptivities of W(VI), ϵ_0 , and WO₃L³⁻, ϵ_1 , at pH 7.5, Different Wavelengths, and Various Ionic Strengths, I , of NaClO₄

$l/\text{mol dm}^{-3}$	ϵ_0			ϵ_1		
	260 nm	265 nm	270 nm	260 nm	265 nm	270 nm
0.1	660.8	486.6	486.6	1116.6	1116.6	1116.6
0.3	527.5	365.0	246.1	1221.6	1268.3	1250.0
0.5	505.8	345.0	245.8	1533.3	1583.3	1716.6
0.7	478.3	320.0	224.1	716.6	750.0	766.6
1.0	522.5	355.0	251.6	2316.6	2350.0	2333.3

the concentration of the complex is

$$[C] = A_c/\epsilon_1 \quad (5)$$

It is now certain that the WO₃L³⁻ ion is tetrahedral in aqueous solution.²⁷ In the usual potentiometric method for evaluating metal–ligand stability constants, the competition between the metal ion and the hydrogen ion for the ligand is studied, and the pH region of interest is from about 1 to 5.²¹ In the W(VI) systems, however, the complication of metal polymerization is introduced in acidic solutions. Because the polymerization equilibria are not well understood, this pH region is not useful for stability determinations. In more alkaline solutions, on the other hand, a pH-dependent process involving the competition between tungstate formation and metal–ligand complexation, i.e., a competition between OH⁻ and ligand for the metal ion, can be utilized. This process was determined from the NMR studies to be important from about pH 6 to 9 and can be represented by²¹



where M represents W and L represents the aminopolycarboxylic acid ligand. In the pH region above 6, no evidence was found for any Mo species containing fewer than three oxygen atoms, e.g., MoO₂2⁺, as has been proposed for other systems.²⁸ The molybdenum-coordinating species in all the aminopolycarboxylic acid systems above pH 6 is MoO₃, and by analogy, we have assumed that the corresponding coordinating unit in the tungsten systems is WO₃.

By use of the potentiometric technique, results obtained for simple one to one metal–ligand chelates are more precise than for higher complexes, such as those formed with EDTA. In fact, for the multicomplex systems, the NMR data are probably more reliable than the potentiometric data as indicated by the range of calculated values for the EDTA systems.²¹ A comparison of the formation constants shows that within experimental error there is essentially no difference between the stabilities of the


Table 5. Average Values of β_{112} at pH 7.5 and Different Ionic Strengths for the Complexation of Tungsten(VI) with NTA, $T = 25^\circ\text{C}$

$l/\text{mol dm}^{-3}$	$\log \beta_{112}$	experimental conditions	ref
0.1	19.00 ± 0.30		this work
0.3	19.37 ± 0.10		this work
0.5	19.65 ± 0.25		this work
0.7	20.10 ± 0.10		this work
1.0	19.40 ± 0.20		this work
	18.86 ± 0.05	$I = 0.15 \text{ M}, T = 25^\circ\text{C}$	21
	19.10 ± 0.2	$I = 1.0\text{--}2.5 \text{ M}, T = 35^\circ\text{C}$	21
	17.75	$I = 0.5 \text{ M NaClO}_4, T = 25^\circ\text{C}$	22
	19.03 ± 0.15	$I = 3 \text{ M NaClO}_4, T = 25^\circ\text{C}$	23

corresponding Mo(VI) and W(VI) chelates. This does not seem too surprising in view of the similarities of the two ions and their nearly equal ionic radii.²¹ However, NMR studies show that tungsten chelates are significantly more labile with respect to individual metal–ligand bonds than are the molybdenum chelates. The slowness with which pH equilibrium was attained in the NTA chelates may be explained by the higher negative charge of this complex compared to the other complexes and the subsequently slower reaction with OH⁻



For the NTA ligand system, NMR studies also indicate that only one metal–ligand species exists above pH 6, MoO₃L³⁻.²¹ Thus, W(VI) will bond with this tridentate ligand as a 1:1 complex



with the stability constant, β_{112} , as

$$\beta_{112} = [\text{WO}_3\text{L}^{3-}]/[\text{WO}_4^{2-}][\text{L}^{3-}][\text{H}^+]^2 \quad (9)$$

The following equations are valid for the total concentration of tungsten (C_M) and the total concentration of the ligand (C_L) at the maximum point on the plot of Figure 1

$$C_M = [W] + [C] \quad (10)$$

$$C_L = [L] + [C] \quad (11)$$


[C] is the concentration of the complex. By substitution of eqs 5, 10, and 11 in eq 9, we can calculate the values of β_{112} according to reaction 8. Stability constants have been calculated by the combination of spectrophotometric and potentiometric data. The values of β_{112} at different ionic strengths together with the values of literature are shown in Table 5.

The dependence of the dissociation and stability constants on the ionic strength can be described according to the previous works^{14–17,25}

$$\log \beta_{112}(I) =$$

$$\log \beta_{112}(I_1) - AZ^* \left(\frac{I^{0.5}}{1 + BI^{0.5}} - \frac{I_1^{0.5}}{1 + BI_1^{0.5}} \right) + C(I - I_1) + D(I^{1.5} - I_1^{1.5}) \quad (12)$$

where I and I_1 are the actual and reference ionic strengths, respectively, and according to eq 13






Figure 2. Plot of $\log K_5$ for WO_3L^{3-} vs the square root of ionic strength.

Table 6. Parameters for the Dependence on the Ionic Strength of Dissociation and Stability Constants at 25 °C

species	C	D	Z
K_5	1.057	-0.864	6
K_2	0.803	-0.730	4
K_1	0.210	-0.775	2
WO_3L^{3-}	7.130	-5.465	6

Table 7. Calculated Stability Constants at pH 7.5 and Different Ionic Strengths for the Complexation of Tungsten(VI) with NTA, $T = 25$ °C

$I/\text{mol dm}^{-3}$	$\log \beta_{112}$
0.1	18.94
0.3	19.38
0.5	19.65
0.7	19.72
1.0	19.51

$Z^* = pm^2 + qr^2 + r - (pm + qn + r)^2$, where m and n are the charges on the metal ion and the ligand, respectively. Considering $A = 0.5115$ and $B = 1.489$, eq 12 can be simplified

$$\log \beta_{112}(I) = \log \beta_{112}(I_1) - Z^* \left(\frac{\rho.5}{2 + 3\rho.5} - \frac{I_1^{0.5}}{2 + 3I_1^{0.5}} \right) + C(I - I_1) + D(I^{1.5} - I_1^{1.5}) \quad (14)$$

where C and D are empirical coefficients and their values were obtained by minimizing the error squares sum, (I_1), and the Gauss-Newton nonlinear least-squares method in a suitable computer program

$$U = \sum_{j=1}^n (a_j - b_j)^2 \quad (j = 1, 2, 3, \dots) \quad (15)$$

where a_j is a quasixperimental quantity and b_j is a calculated one. The values of C and D are shown in Table 6. In this research, a_j is the experimental stability constant and b_j is the calculated one. We have used $I_1 = 0.5$ as the reference ionic strength in order to obtain better consistency between experimental and calculated stability constants. The calculated stability constants are given in Table 7.

The dependence on ionic strength of $\log \beta_{112}$ is shown in Table 4, which is in good agreement with the results obtained in the previous papers.^{14-17,25} The dependence of β_{112} on the ionic strength is shown in Figure 2. This figure shows that β_{112} has one maximum at $I = 0.7$ similar to the maximum for the $\text{Mo(VI)}-\text{NTA}$ system in our previous paper.²⁵

Literature Cited

- Raajmakers, H.; Macielra, S.; Dias, J. M.; Teixeira, S.; Burasakov, S.; Huber, R.; Moura, J. J. G.; Moura, I.; Romo, M. J. Gene Sequence and the 1.8 Crystal Structure of the Tungsten-Containing Formate Dehydrogenase from *Desulfovibrio gigas*. *Structure* 2002, 10, 1261-1272.
- Lorber, C.; Donahue, J. P.; Goddard, C. A.; Nord, E.; Holm, R. H. Synthesis, Structures, and Oxo Transfer Reactivity of Bis(dithiolene) Tungsten(IV, VI) Complexes Related to the Active Sites of Tungstoenzymes. *J. Am. Chem. Soc.* 1998, 120, 8102-8112.
- Gielzak, K.; Wojciechowski, W. Complexes of Tungsten(VI) with Tryptophan. *Pol. J. Chem.* 1988, 62, 559-562.
- Yang, C. H.; Dzugan, S. J.; Goedken, V. L. Structure and Reactivity of an Unusual Tungsten(VI) Porphyrin Complexes having Cis Oxo and Peroxo Groups. *J. Chem. Soc., Chem. Commun.* 1985, 20, 1425-1426.
- Daniele, P. G.; Rigano, C.; Sammartano, S. Ionic Strength Dependence of Formation Constants, Protonation Constants of Organic and Inorganic Acids. *Talanta* 1983, 30, 81-87.
- De Stefano, C.; Foti, C.; Gianguzza, A. Salt Effects on the Protonation and on Alkali and Alkaline Earth Metal Complex Formation of 1,2,3-Propanetricarboxylate in Aqueous Solution. *Talanta* 1994, 41, 1715-1722.
- Daniele, P. G.; Rigano, C.; Sammartano, S.; Zelano, V. Ionic Strength Dependence of Formation Constants XVIII. The Hydrolysis of Iron (III) in Aqueous KNO_3 Solutions. *Talanta* 1994, 41, 1577-1582.
- Gil, R.; Corbillon, M. S.; Olazabal, M. A.; Madariaga, J. M.; Potentiometric Study of the Protonation Equilibrium of Tris-(Hydroxymethyl) Aminoethane in Aqueous Sodium Perchlorate Solutions at 25 °C: Construction of a Thermodynamic Model. *Talanta* 1997, 44, 891-896.
- Daniele, P. G.; Rigano, C.; Sammartano, S. Ionic Strength Dependence of Formation Constants. Alkali Metal Complexes of EDTA, NTA, Diphosphate, and Triphosphosphate in Aqueous Solution. *Anal. Chem.* 1985, 57, 2056-2060.
- De Robertis, A.; Foti, C.; Gianguzza, A. Ionic Strength Dependence of Formation Constants. Salt Effects on the Protonation of 1H-1,2,3-Triazole-4,5-dicarboxylic Acid in Aqueous Solution. *J. Chem. Res.* 1995, 288-289.
- De Robertis, A.; De Stefano, C.; Sammartano, S.; Rigano, C. The Determination of Formation Constants of Weak Complexes by Potentiometric Measurements: Experimental Procedures and Calculation Methods. *Talanta* 1987, 34, 933-938.
- De Stefano, C.; Gianguzza, A.; Piazzese, D.; Sammartano, S. Polyacrylate Protonation in Various Aqueous Ionic Media at Different Temperatures and Ionic Strengths. *J. Chem. Eng. Data* 2000, 45, 876-881.
- Daniele, P. G.; De Robertis, A.; De Stefano, C.; Sammartano, S.; Rigano, C. On the Possibility of Determining the Thermodynamic Parameters for the Formation of Weak Complexes Using a Simple Model for the Dependence on Ionic Strength of Activity Coefficients: Na^+ , K^+ and Ca^{2+} Complexes of Low Molecular Weight Ligands in Aqueous Solution. *J. Chem. Soc., Dalton Trans.* 1985, 2355-2361.
- Charif, F.; Afrazeh Dogaheh, L. Ionic Strength Dependence of Formation Constants: Complexation of Cysteine with Molybdenum(VI). *J. Chem. Eng. Data* 2003, 48, 999-1003.
- Majlesi, K.; Zare, K. Comparison of the Ionic Strength Effect on the Complexation of Dioxovanadium(V) with Phenylalanine and Molybdenum(VI) with Glutamic Acid. Proceedings of 6th Iranian Physical Chemistry Seminar, Faculty of Science, Urmia University, Urmia, Iran August 27-29, 2002.
- Charif, F.; Zare, K.; Majlesi, K. Ionic Strength Dependence of Formation Constants: Complexation of Molybdenum(VI) with Glutamic Acid. *J. Chem. Eng. Data* 2000, 45, 833-836.
- Charif, F.; Zare, K.; Majlesi, K. Ionic Strength Dependence of Formation Constants, Protonation and Complexation of Phenylalanine with Dioxovanadium(V) at Different Temperatures. *J. Chem. Res.* 2000, 186-187.
- Marcu, G.; Tomus, M. Radiochromatographic and Electrochromatographic Study of Behavior of Sodium Tungstate Solutions Under the Action of Nitrotriacetic Acid. *Ser. Chem. 1968*, 13, 11-14.
- Kula, R. J. Solution Equilibria and Structures of Molybdenum(VI) Chelates, *N*-Methyliminodiacetic Acid. *Anal. Chem.* 1966, 38, 1382-1388.
- Kula, R. J. Solution Equilibria and Structures of Molybdenum(VI) Chelates, (Ethylendinitrilo) Tetracetic Acid. *Anal. Chem.* 1966, 38, 1581-1584.
- Kula, R. J.; Rabenstein, D. L. Potentiometric Determination of Stabilities of Molybdenum(VI) and Tungsten(VI) Chelates. *Anal. Chem.* 1966, 38, 1934-1936.

- (22) Collin, J. P.; Lagrange, P. *Bull. Soc. Chim. France*, 1976, 9–10, 1304.
- (23) Zare, K.; Lagrange, P.; Lagrange, J. Determination and Comparison of Stability Constants of Vanadium(V), Molybdenum(VI) and Tungsten(VI) Aminocarboxylate Complexes. *J. Chem. Soc., Dalton Trans.* 1979, 1372–1376.
- (24) Anderegg, G. Critical Survey of Stability Constants of NTA Complexes. *Pure Appl. Chem.* 1982, 54, 2693–2758.
- (25) Majlesi, K.; Zare, K.; Teimouri, F. Ionic Strength Dependence of Formation Constants, Protonation and Complexation of Nitrotriacetic Acid with Molybdenum(VI). *J. Chem. Eng. Data* 2003, 48, 680–683.
- (26) Billo, Joseph, E. *Excel for Chemists*, 2nd ed.; John Wiley & Sons: New York, 2001.
- (27) Cotton, F. A.; Wilkinson, G. *Advanced Inorganic Chemistry*, 3rd ed.; Wiley-Interscience: New York, 1972.
- (28) Spengler, G.; Ganshelmer, J. *Angew. Chem.* 1957, 69, 523.
- (29) Dean, J. A.; *Langes Handbook of Chemistry*, 14th ed.; McGraw-Hill: New York, 1992.

Received for review May 15, 2003. Accepted December 15, 2003.

JE034092U